

THE RELATIONSHIP OF PASSIVE
SMOKING TO VARIOUS HEALTH
OUTCOMES AMONG SEVENTH DAY
ADVENTISTS IN CALIFORNIA

BUTLER, TERRENCE LADLIT
DEGREE DATE: 1998

UNIVERSITY OF CALIFORNIA
SAN DIEGO

2023513023

This is an authorized facsimile, made from the microfilm master copy of the original dissertation or masters thesis published by UMI.

The bibliographic information for this thesis is contained in UMI's Dissertation Abstracts database, the only central source for accessing almost every doctoral dissertation accepted in North America since 1861.

U·M·I Dissertation
Information Service

University Microfilms International
A Bell & Howell Information Company
300 N. Zeeb Road, Ann Arbor, Michigan 48106
800-521-0600 OR 313-761-4700

Printed in 1991 by xerographic process
on acid-free paper

3266

2023513024

Order Number 8827026

The relationship of passive smoking to various health outcomes
among Seventh-day Adventists in California

Butler, Terrence Leslie, Dr.P.H.

University of California, Los Angeles, 1988

Copyright ©1988 by Butler, Terrence Leslie. All rights reserved.

U·M·I
300 N. Zeeb Rd.
Ann Arbor, MI 48106

2023513025

INFORMATION TO USERS

The most advanced technology has been used to photograph and reproduce this manuscript from the microfilm master. UMI films the text directly from the original or copy submitted. Thus, some thesis and dissertation copies are in typewriter face, while others may be from any type of computer printer.

The quality of this reproduction is dependent upon the quality of the copy submitted. Broken or indistinct print, colored or poor quality illustrations and photographs, print bleedthrough, substandard margins, and improper alignment can adversely affect reproduction.

In the unlikely event that the author did not send UMI a complete manuscript and there are missing pages, these will be noted. Also, if unauthorized copyright material had to be removed, a note will indicate the deletion.

Oversize materials (e.g., maps, drawings, charts) are reproduced by sectioning the original, beginning at the upper left-hand corner and continuing from left to right in equal sections with small overlaps. Each original is also photographed in one exposure and is included in reduced form at the back of the book. These are also available as one exposure on a standard 35mm slide or as a 17" x 23" black and white photographic print for an additional charge.

Photographs included in the original manuscript have been reproduced xerographically in this copy. Higher quality 6" x 9" black and white photographic prints are available for any photographs or illustrations appearing in this copy for an additional charge. Contact UMI directly to order.

U·M·I

University Microfilms International
A Bell & Howell Information Company
300 North Zeeb Road, Ann Arbor, MI 48106-1346 USA
313 761-4700 800 521-0600

2023513026

UNIVERSITY OF CALIFORNIA

Los Angeles

The Relationship of Passive Smoking to Various
Health Outcomes among Seventh-day Adventists in California

A dissertation submitted in partial satisfaction
of the requirements for the degree
Doctor of Public Health


by


Terrence Leslie Butler


1988


2023513027

The dissertation of Terrence Leslie Butler is approved.


Sander Greenland


William Shonick


Henry Gong


Murray Jarnik


Barbara Visscher, Committee Chair

University of California, Los Angeles

1988

2023513028

© Copyright by
Terrence Leslie Butler

1988

2023513029

Dedicated to the memory of my mother

Sarah Joan Butler (1922-1987)

A life of immeasurable qualities

2023513030

TABLE OF CONTENTS

	Page
List of Figures	vii
List of Tables	viii
Acknowledgements	xvi
Vita, Publications and Presentations	xx
Abstract	xxi
Chapter 1 INTRODUCTION	1
1.1 Background	1
1.2 Research Hypotheses	3
1.3 Specific Objectives	4
1.4 Definitions and Classification of Terms	5
1.5 Significance, Relevance and Rationale	8
1.6 Organization of Thesis	9
 Chapter 2 LITERATURE REVIEW	 11
2.1 Introduction	11
2.2 Comparison of Mainstream and Sidestream Smoke	13
2.3 Measurement of Passive Smoking Exposure	15
2.4 General Health Effects	20
2.5 Plausibility of a Health Effect	22
2.6 Lung Cancer	23

	Page
2.7 Other Cancers	32
2.8 Heart Disease	36
2.9 All Cause Mortality	40
2.10 Summary	40
Chapter 3 METHODS	43
3.1 Background of Study	43
3.2 Study Design	52
3.3 Description of the Study Populations	52
3.4 Data Description	58
3.5 Outcome Ascertainment Procedures	61
3.6 Passive Smoking Exposure Assessment	65
3.7 Definitions of Other Exposure Variables	70
3.8 Data Analysis and Statistical Methods	76
3.9 Limitations and Potential Problems of Study	79
Chapter 4 DESCRIPTIVE STATISTICS	81
4.1 Spouse Pairs Demographic Data	81
4.2 Smoking Exposure	82
4.3 AHSMOG Cohort Demographic Data	90

2023513032

	Page
Chapter 5 LUNG CANCER RESULTS	97
5.1 Spouse Pairs Cohort	97
5.2 AHSMOG Cohort	105
5.3 Summary	112
Chapter 6 SMOKING RELATED CANCER RESULTS	114
6.1 Spouse Pairs Cohort	114
6.2 AHSMOG Cohort	131
6.3 Summary	135
Chapter 7 ALL INCIDENT CANCER RESULTS	140
7.1 Spouse Pairs Cohort Incident Cancers	140
7.2 AHSMOG Cohort All Incident Cancers	145
7.3 Summary	149
Chapter 8 MORTALITY RESULTS	153
8.1 Spouse Pairs Cohort - All Cause Mortality Results	153
8.2 Spouse Pairs Cohort - Ischemic Heart Disease Mortality Results	160
8.3 AHSMOG Cohort - All Cause Mortality Results	163
8.4 AHSMOG Cohort - Ischemic Heart Disease Mortality Results	173

	Page
8.5 Summary	176
Chapter 9 DISCUSSION AND CONCLUSIONS	180
9.1 Data Quality	180
9.2 Lung Cancer	187
9.3 Smoking Related Cancers	191
9.4 Cervical Cancer	194
9.5 All Incident Cancers	196
9.6 Mortality Results	198
9.7 Conclusions	199
References	203
Appendices 1 Glossary of Terminology and Abbreviations	214
2 AHS Census Questionnaire (1974)	216
3 AHS Lifestyle Questionnaire (1976)	226
4 AHSMOG Questionnaire (1977)	248
5 AHSMOG Questionnaire (1987)	254
6 Data Collection Forms	269
1977 Hospital History Form	270
1983 Hospital History Form	271
Discharge Diagnosis Form	273
Tumor Abstract Form	275

LIST OF FIGURES

	Page
Figure 3.1 Flow Chart of Adventist Health Study	47
3.2 Relationship of Spouse Pairs and AHSMOG Sub-Populations to the Adventist Health Study Incidence Population	53

LIST OF TABLES

Table 1.1 Smoking-Related Cancers	7
2.1 Distribution of Constituents in Mainstream Smoke (MS) and the Ratio of Sidestream Smoke (SS) to MS of non-filter cigarettes	14
2.2 Summary of Case Control Studies of Lung Cancer	25
2.3 Summary of Cohort Studies of Lung Cancer	31
2.4 Summary of Studies of Other Cancers	33
2.5 Summary of Studies of Heart Disease	37
2.6 Summary of Studies of All Cause Mortality	41
3.1 Age-Standardized Mortality Ratios for Selected Causes of Death Among 25,000 White California Adventists, 1960-1980	45
3.2 Base Line Variables Measured on the Three Self-Administered Questionnaires, 1974-1977	50
3.3 Spouse Pairs Cohort: Selected Descriptive Data	55
3.4 AHSMOG Cohort: Selected Descriptive Data	57
3.5 Spouse Pairs Cohort: Spouse Smoking Status in Marriage	68

		Page
Table 3.6	Percentage Agreement Between Two Methods of Classifying Husband's Smoking Status in Marriage	70
4.1	Spouse Pairs Cohort: Sex-Specific Distribution of Age at LO by Age Groups	81
4.2	Spouse Pairs Cohort: Sex-Specific Distribution of Smoking Status, 1976	82
4.3	Spouse Pairs Cohort: Sex-Specific Distribution of Number of Cigarettes Regularly Smoked	83
4.4	Spouse Pairs Cohort: Sex-Specific Distribution of Number of Years Regularly Smoked	84
4.5	Spouse Pairs Cohort: Sex-Specific Distribution of Pack Years of Cigarettes Smoked	85
4.6	Spouse Pairs Cohort: Smoking Status of Spouse Pairs	86
4.7	Spouse Pairs Cohort: Spouse Smoking Status in Marriage	87
4.8	Spouse Pairs Cohort: Sex-Specific Mortality, 1976-1982, by Underlying Causes	88
4.9	Spouse Pairs Cohort: Sex and Site Specific Incident Cases of Cancer, 1977-1982	89
4.10	AHSMOG Cohort: Sex-Specific Distribution of Age Groups	90
4.11	AHSMOG Cohort: Sex-Specific Distribution of Smoking Status	91
4.12	AHSMOG Cohort: Sex-Specific Distribution of Cigarettes Smoked	91
4.13	AHSMOG Cohort: Sex-Specific Distribution of Years Smoked	92

		Page
Table 4.14	AHSMOG Cohort: Sex-Specific Distribution of Pack Years of Cigarettes Smoked	92
4.15	AHSMOG Cohort: Sex-Specific Distribution of Years Lived with a Smoker	93
4.16	AHSMOG Cohort: Sex-Specific Distribution of Years Worked with a Smoker	94
4.17	AHSMOG Cohort: Sex-Specific Mortality, 1976-1982, by Underlying Causes	95
4.18	AHSMOG Cohort: Sex and Site Specific Incident Cases of Cancer, 1977-1982	96
5.1	Spouse Pairs Females: Histology of Lung Cancer Cases	98
5.2	Spouse Pairs Females: Lung Cancer, 1977-1982, Distribution of Incident Cases, Person Years, and Incidence Rates by Age Group and Husbands Smoking Status in Marriage	99
5.3	Spouse Pairs Females: Age Adjusted Rate Ratios of Lung Cancer, 1977-1982, for Selected Exposure Factors	101
5.4	Spouse Pairs Female Non-Smokers: Age Adjusted Rate Ratio for Lung Cancer, 1977-1982, for Selected Exposure Factors	104
5.5	Spouse Pairs Female Non-Smokers: Comparison of Lung Cancer, Age Adjusted Rate Ratios for Exposure to Spouse Smoking using Different Statistical Methods	105
5.6	AHSMOG Cohort: Histology of Lung Cancer Cases	106
5.7	AHSMOG Current Non-Smokers: Lung Cancer, 1977-1982, Distribution of Incident Cases, Person Years and Incidence Rates by Age Group and Years Lived with a Smoker	107

		Page
Table 5.8	AHSMOG Current Non-Smokers: Lung Cancer, 1977-1982, Distribution of Incident Cases, Person Years and Incidence Rates by Age Group and Years Worked with a Smoker	108
5.9	AHSMOG Females: Adjusted Rate Ratios of Lung Cancer, 1977-1982, for Selected Exposure Factors	110
5.10	AHSMOG Males: Adjusted Rate Ratios of Lung Cancer, 1977-1982, For Selected Exposure Factors	111
5.11	AHSMOG Lung Cancer Adjusted Rate Ratios for Varying Years and Types of Passive Smoking Exposure	113
6.1	Spouse Pairs Females: Site Specific Distribution of Smoking Related Cancers	115
6.2	Spouse Pairs Females: All Smoking Related Cancers, 1977-1982, Distribution of Incident Cases, Person Years and Incidence Cases by Age Group and Husband's Smoking Status in Marriage	116
6.3	Spouse Pairs Females: Age Adjusted Rate Ratio of Smoking Related Cancer Excluding Lung Cancer for Selected Exposure Factors, 1977-1982	117
6.4	Spouse Pairs Females: Age Adjusted Rate Ratio of All Smoking Related Cancer for Selected Exposure Factors	119
6.5	Spouse Pairs Female Non-Smokers: Age Adjusted Rate Ratio of Smoking Related Cancer Excluding Lung for Selected Exposure Factors, 1977-1982	121
6.6	Spouse Pairs Female Non-Smokers: Age Adjusted Rate Ratio of Smoking Related Cancer Including Lung, 1977-1982, for Selected Exposure Factors	122
6.7	Spouse Pairs Female Non-Smokers: Smoking Related Cancers Age Adjusted Rate Ratios for Females Married to a Smoker or Married to a Non-Smoker	123

x

2023513038

		Page
Table 6.8	Spouse Pairs Females: Cervical Cancer, 1977-1982. Distribution of Incident Cases, Person Years and Incidence Rates by Age Group and Husband Smoking Status in Marriage	124
6.9	Spouse Pairs Females: Age Adjusted Rate Ratio of Cervical Cancer, 1977-1982, for Selected Exposure Factors	126
6.10	Spouse Pairs Female Non-Smokers: Age Adjusted Rate Ratio of Cervical Cancer, 1977-1982, for Selected Exposure Factors	128
6.11	Spouse Pairs Female Non-Smokers: Comparison of Adjusted Rate Ratios of Cervical Cancer, 1977-1982, when Stratified on Different Risk Factors	129
6.12	Spouse Pairs Female Non-Smokers: Cervical Cancer Age Adjusted Rate Ratios for Females Married to a Smoker or Married to a Non-Smoker	130
6.13	AHSMOG Cohort: Site Specific Distribution of Smoking Related Cancers, 1977-1982	131
6.14	AHSMOG Current Non-Smokers: All Smoking Related Cancers, 1977-1982, Distribution of Incident Cases, Person Years and Incidence Rates by Age Group and Years Lived with a Smoker	133
6.15	AHSMOG Current Non-Smokers: All Smoking Related Cancers, 1977-1982, Distribution of Incident Cases, Person Years and Incidence Rates by Age Group and Years Worked with a Smoker	134
6.16	AHSMOG Females: Adjusted Rate Ratios of All Smoking Related Cancer, 1977-1982, for Selected Exposure Factors	137
6.17	AHSMOG Males: Adjusted Rate Ratios of All Smoking Related Cancer, 1977-1982, for Selected Exposure Factors	138

		Page
Table 6.18	AHSMOG: All Smoking Related Cancers Age Adjusted Rate Ratio for Varying Years and Types of Passive Smoking Exposure	138
7.1	Spouse Pairs Females: All Incident Cancers, 1977-1982, Distribution of Incident Cases, Person Years and Incidence Rates by Age Group and Husband Smoking Status in Marriage	141
7.2	Spouse Pairs Females: Age Adjusted Rate Ratio of All Incident Cancer for Selected Exposure Factors, 1977-1982	143
7.3	Spouse Pairs Non-Smoking Females: Age Adjusted Rate Ratio of All Incident Cancer, 1977-1982, for Selected Exposure Factors	144
7.4	Spouse Pairs Females: Adjusted Rate Ratio of All Incident Cancers, 1977-1982, for Selected Exposure Factors	146
7.5	AHSMOG Current Non-Smokers: All Incident Cancers, 1977-1982, Distribution of Incident Cases, Person Years and Incidence Rates by Age Group and Years Lived with a Smoker	147
7.6	AHSMOG Current Non-Smokers: All Incident Cancers, 1977-1982, Distribution of Incident Cases, Person Years and Incidence Rates by Age Group and Years Worked with a Smoker	148
7.7	AHSMOG Females: Adjusted Rate Ratio of All Incident Cancers, 1977-1982, for Selected Exposure Factors	150
7.8	AHSMOG Males: Adjusted Rate Ratio of All Incident Cancers, 1977-1982, for Selected Exposure Factors	151
8.1	Spouse Pairs Females: All Cause Mortality, 1977-1982, Distribution of Deaths, Person Years and Incidence Rates by Age Group and Husband Smoking Status in Marriage	154

218

2023513040

		Page
Table 8.2	Spouse Pairs Females: Age Adjusted Rate Ratio of All Natural Cause Mortality, 1976-1982, for Selected Exposure Factors	156
8.3	Spouse Pairs Non-Smoking Females: Age Adjusted Rate Ratio of All Natural Cause Mortality, 1976-1982, for Selected Exposure Factors	158
8.4	Spouse Pairs Females: Ischemic Heart Disease Mortality, 1976-1982, Distribution of Deaths, Person Years and Incidence Rates by Age Group and Husband Smoking Status in Marriage	161
8.5	Spouse Pairs Non-Smoking Females: Age Adjusted Rate Ratio of Ischemic Heart Disease Mortality, 1976-1982, for Selected Exposure Factors	162
8.6	AHSMOG Current Non-Smokers: All Cause Mortality, 1976-1982, Distribution of Deaths, Person Years and Incidence Rates by Age Group and Years Lived with a Smoker	164
8.7	AHSMOG Current Non-Smokers: All Cause Mortality, 1976-1982, Distribution of Deaths, Person Years and Incidence Rates by Age Group and Years Worked with a Smoker	165
8.8	AHSMOG Females: Age Adjusted Rate Ratio of All Cause Mortality, 1976-1982, for Selected Exposure Factors	167
8.9	AHSMOG Males: Age Adjusted Rate Ratio of All Cause Mortality, 1976-1982, for Selected Exposure Factors	169
8.10	AHSMOG Current Non-Smokers: Ischemic Heart Disease Mortality, 1976-1982, Distribution of Deaths, Person Years and Incidence Rates by Age Group and Years Lived with a Smoker	171

		Page
Table 8.11	AHSMOG Current Non-Smokers: Ischemic Heart Disease Mortality, 1976-1982, Distribution of Deaths, Person Years and Incidence Rates by Age Group and Years Worked with a Smoker	172
8.12	AHSMOG Females: Age Adjusted Rate Ratio of Ischemic Heart Disease Mortality, 1976-1982, for Selected Exposure Factors	174
8.13	AHSMOG Males: Age Adjusted Rate Ratio of Ischemic Heart Disease Mortality, 1976-1982, for Selected Exposure Factors	177
9.1	Spouse Pairs: Subject Smoking Status Percent Agreement Between 1976 and 1987 Responses	182
9.2	Spouse Pairs Females: Percent Agreement Between Created Variable "Husband Smoked in Marriage" and 1987 Response "Lived with Husband who Smoked"	184
9.3	AHSMOG Cohort: Agreement Between 1977 and 1987 Responses to ETS Exposure Questions	186
9.4	A Comparison of Relative Risks of Lung Cancer for ETS Exposure among the AHSMOG and Spouse Pairs Cohorts	188
9.5	A Comparison of Relative Risks of All Smoking Related Cancer for ETS Exposure among the AHSMOG and Spouse Pairs Cohorts	193
9.6	A Comparison of Relative Risks of All Incident Cancers for ETS Exposure among the AHSMOG and Spouse Pairs Cohorts	197
9.7	A Comparison of Relative Risks of All Cause Mortality for ETS Exposure among the AHSMOG and Spouse Pairs Cohorts	201
9.8	A Comparison of Relative Risks of Ischemic Heart Disease for ETS Exposure among the AHSMOG and Spouse Pairs Cohorts	202

CHAPTER 5: LUNG CANCER RESULTS

In this chapter the results of the lung cancer analyses are presented for both the spouse pairs females and the AHSMOG cohort. For the spouse pairs females the initial analyses include both the current and past smokers. However, later analyses are restricted to the non-smoking population of this cohort. Although the major variables of interest are the passive smoking exposures, other selected factors are chosen for inclusion in the analyses because they may be independent risk factors or have some protective effect on the outcome. The methods of analyses included the calculation of a crude measure of effect and stratified analyses.

5.1 SPOUSE PAIRS COHORT

In the female population of the spouse pairs cohort nine incident cases of primary lung cancer were diagnosed during the follow-up period 1977 to 1982. Histological confirmation was obtained for each case and the information abstracted from medical pathology reports. The distribution by histological type is presented in Table 5.1. Since there were very few cases these were grouped together for analytical purposes and no effort was made to differentiate by various histological types.

TABLE 5.1
SPOUSE PAIRS - FEMALES
HISTOLOGY OF LUNG CANCER CASES

Histological Type	Number
Large cell carcinoma	1
Old cell carcinoma	1
Adenocarcinoma	7

All the cases occurred in the age range of 45 to 69 and only one of the nine cases was diagnosed in a former smoker. No cases occurred among females classified as current smokers while eight cases were among non-smokers. Table 5.2 presents the distribution of incident cases, person-years and incidence rates by ten year age groups according to the smoking status of subject and spouse.

A stratified analysis, stratifying on ten year age groups, was performed and Mantel-Haenszel summary rate ratios and corresponding 95% confidence intervals were calculated for selected exposure factors. The incidence rates, crude rate ratios, age adjusted rate ratios, confidence intervals and p-values are shown in Table 5.3. The differences between the crude rate ratios and adjusted rate ratios indicate that there was a

2023513045

Age Group	HUSBAND SMOKER		HUSBAND NON-SMOKER		WIFE SMOKER		WIFE NON-SMOKER	
	Cases/ Rate/ Person years	10,000 PY	Cases/ Rate/ Person years	10,000 PY	Cases/ Rate/ Person years	10,000 PY	Cases/ Rate/ Person years	10,000 PY
25-34	0	942	0	8004	0	718	0	836
35-44	0	1552	0	8637	0	899	0	816
45-54	1	2190	2	9520	0	1011	0	842
55-64	1	3000	2	8578	0	1240	0	876
65-74	1	2159	1	5980	0	629	0	305
75-84	0	682	0	2112	0	70	0	39
85-94	0	63	0	220	0	0	0	8
Total	3	10578	2	84	6	43052	1	4668
							2.14	0.3619
								0.00

TABLE 5.2
SPOUSE PAIRS - FEMALES
LUNG CANCER (1977-1982)
DISTRIBUTION OF INCIDENT CASES, PERSON YEARS AND INCIDENCE RATES BY AGE GROUP
AND HUSBAND SMOKING STATUS IN MARRIAGE

confounding effect by age that to some extent was accounted for in the stratified analysis. The small number of cases and low statistical power militate against the possibility of achieving statistically significant results. It is also impossible to assess effect modification with so few cases.

For our major exposure factor of interest, husband's smoking status in marriage, summary rate ratios of 1.94 (95% C.I. 0.46-8.24) and 2.47 (95% C.I. 0.29-21.18) were obtained for past and current exposure respectively when compared to the referent group of never exposed. A similar doubling of the risk was observed when the husband's smoking status was dichotomized into never and ever smoked, RR of 2.04 (95% C.I. 0.54-7.65). No increased risk of lung cancer was observed in this population for the subjects own smoking status. However, when smokers of more than ten pack years of cigarette smoking were compared to the referent group of non-smokers there was increased risk observed, RR of 2.22 (95% C.I. 0.28-17.74). The solitary case in the exposed category illustrates the equivocal nature of these results.

There is a suggestion of a decreased risk for those subjects who have attended college versus those with a high school education or less and an increased risk for subjects with blue collar working spouses compared

TABLE 53

SPOUSE PAIRS - FEMALES
AGE ADJUSTED RATE RATIOS (RR) OF LUNG CANCER (1977-1982)
FOR SELECTED EXPOSURE FACTORS

Factor	Levels	* Cases/ Person Years	* Rate/ 10 ⁶ PY	Crude RR	Adjusted RR ^a (95% C.I.)	
Husband Smoking status in marriage	Never Past Current	6 47278 3 11564 1 3912	106 259 256	Ref. 2.44 2.41	Ref. 1.94 (0.46-8.24) 2.47 (0.29-21.18)	p = 0.53
Husband smoked in marriage	No Yes	6 47278 4 15560	106 257	Ref. 2.43	Ref. 2.04 (0.54-7.65)	p = 0.27
Female smoking status	Never Past Current	8 54891 1 7559 0 884	148 132 0	Ref. 0.91 0.00	Ref. 0.86 (0.11-6.92) 0.00	p = 0.94
Female Smoked	Non-smoker Smoker	8 54891 1 8443	148 118	Ref. 0.81	Ref. 0.80 (0.10-6.38)	p = 0.83
Pack years of smoking female	None 0-10 > 10	8 56772 0 3534 1 2827	141 0 354	Ref. 0.00 2.51	Ref. 0.00 2.22 (0.28-17.74)	p = 0.58
Education	High School College +	4 21036 4 43263	190 116	Ref. 0.61	Ref. 0.75 (0.20-2.83)	p = 0.64
Live in rural area	Yes No	3 30201 6 32490	99 154	Ref. 1.55	Ref. 1.59 (0.38-6.65)	p = 0.53

101

2023513047

TABLE 5.3 continued

Factor	Levels	Cases/ Person Years	Rate 10 ⁶ PY	Crude RR	Adjusted RR* (95% CI)
Spouse Occupation	White Collar	2 34665	68	Ref.	Ref.
	Blue Collar	6 26120	230	3.96	4.15 (0.78-22.15) p = 0.10
Fruit Index	11 + times week	4 26189	153	Ref.	Ref.
	4-10 times week	4 23951	167	1.09	1.22 (0.31-4.90) p = 0.87
Beta Carotene Index	0-3 times week	1 12005	83	0.54	0.64 (0.07-5.83) p = 0.87
	High	4 22404	178	Ref.	Ref.
	Medium	4 19882	201	1.12	1.23 (0.31-4.92) p = 0.46
	Low	1 21317	47	0.26	0.31 (0.03-2.82) p = 0.46

* Discrepancies in cases and person years are due to missing values in exposure factors

+ Rate per 1 million person years

- Mantel-Haenszel summary rate ratio (RR) - adjusted for age

Ref. Referent group

Total Subjects - 11,060
Lung Cancer Cases - 8
Total Person Years - 64,487

to subjects with white collar working spouses. However, both results are subject to bias because of the unknown classification of one of the cases and the missing number of person years.

Since there was only one case among the 1,475 females who had ever smoked, it was difficult to assess the influence of active smoking on the overall effect of ETS exposure. Therefore further stratified analyses were restricted to the 9,378 never smoking females. The results of these analyses are presented in Table 5.4. Somewhat similar results are observed as in the previous analysis and the same caveats concerning effect modification, bias and statistical significance apply. For the variable husband smoked in marriage, the age-adjusted rate ratio was 2.02 (95% C.I. 0.48-8.56).

An additional analysis using the conditional maximum likelihood estimator and an exact method for sparse data was performed and the result is compared with the Mantel-Haenszel estimates in Table 5.5. These results have similar point estimates with the mid probability exact binomial confidence intervals being somewhat wider and more conservative.

TABLE 5.4

SPOUSE PAIRS - FEMALE NON-SMOKERS
AGE ADJUSTED RATE RATIOS (RR) FOR LUNG CANCER (1977-1982)
FOR SELECTED EXPOSURE FACTORS

Factor	Levels	* Cases/ Person Years	* Rate/ 10 ⁶ PY	Crude RR	Adjusted RR [†] (95% C.I.)	
Husband Smoking status in marriage	Never	5 43037	116	Ref.	Ref.	
	Past	2 8090	247	2.13	1.69 (0.32-8.88)	
	Current	1 2486	402	3.47	3.37 (0.39-29.05)	p = 0.46
Husband smoked in marriage	No	5 43037	116	Ref.	Ref.	
	Yes	3 10576	284	2.45	2.02 (0.48-8.56)	p = 0.32
Education	High School	3 16063	187	Ref.	Ref.	
	College +	5 38670	129	0.69	0.86 (0.20-3.67)	p = 0.82
Live in rural area	Yes	3 26013	115	Ref.	Ref.	
	No	4 27694	144	1.26	1.30 (0.29-6.81)	p = 0.73
Spouse Occupation	White collar	2 31217	64	Ref.	Ref.	
	Blue collar	5 20768	289	4.51	4.89 (0.89-26.71)	p = 0.06
North Carolina Index	High	4 19454	206	Ref.	Ref.	
	Medium	3 17065	176	0.85	0.93 (0.21-4.17)	
	Low	1 17680	67	0.27	0.33 (0.04-2.99)	p = 0.59

* Discrepancies in cases and person years are due to missing values in exposure factors
 * Rate per 1 million person years
 * Mantel-Haenszel summary rate ratio (RR) - adjusted for age
 Ref. Referent Group

Total Subjects - 9,378
 Lung Cancer Cases - 8
 Total Person Years - 54,894

2023513050

TABLE 5.5
SPOUSE PAIRS - FEMALE NON-SMOKERS
A. COMPARISON OF LUNG CANCER AGE ADJUSTED RATE RATIOS
FOR EXPOSURE TO SPOUSE SMOKING USING
DIFFERENT STATISTICAL METHODS

FACTOR	LEVELS	N	METHOD	ADJUSTED RR (95% C.I.)
Husband smoked in Marriage	No	84007	Mantel-Haenszel	Ref.
	Yes	310575		2.02 (0.48-8.56)
	No		Maximum Likelihood	Ref.
	Yes			2.01 (0.39-8.79)*

* cases/person years
* mid probability (Miettinen); binomial confidence intervals

5.2 AHSMOG COHORT

During the years 1977-1982 thirteen incident cases of lung cancer were diagnosed in the current non-smokers of the AHSMOG cohort. Seven of the cases were males and six were females. The histological types of tumors are presented in Table 5.6. As with the spouse pairs cohort the most predominant histological type was adenocarcinoma. However, all histological types were grouped together for analytical purposes.

TABLE 5.6

AHSMOG
HISTOLOGY OF LUNG CANCER CASES

Histological Type	Male	Female
Large cell carcinoma	1	
Carcinoma NOS	1	1
Squamous cell carcinoma	1	
Adenocarcinoma	4	4
Unknown		1

For females, cases occurred in the age range of 55 to 94 years while for males, cases were limited to the 55-84 year age range. The distribution of incident cases, person years and incidence rates by ten year age groups for the two ETS exposures of interest—years lived with a smoker and years worked with a smoker—are presented in Tables 5.7 and 5.8 respectively. There is an increased risk for age, however, the lack of sufficient cases prevent a careful assessment of trend across passive smoking exposures.

TABLE 5.7

ATISMOO - CURRENT NON SMOKERS
LUNG CANCER (1977-1982)
DISTRIBUTION OF INCIDENT CASES, PERSON YEARS AND INCIDENCE RATES BY AGE GROUP
AND YEARS LIVED WITH A SMOKER

FEMALE Age Group	YEARS LIVED WITH SMOKER					
	NONE		1-10 YEARS		11 + YEARS	
	Cases/ Person years	Rate/ 10,000 PY	Cases/ Person years	Rate/ 10,000 PY	Cases/ Person years	Rate/ 10,000 PY
25-34	0 636	0.00	0 243	0.00	0 398	0.00
35-44	0 1671	0.00	0 420	0.00	0 1149	0.00
45-54	0 2617	0.00	0 847	0.00	0 1628	0.00
55-64	0 3298	0.00	0 872	0.00	1 2269	4.41
65-74	1 2687	3.72	0 677	0.00	1 1791	6.68
75-84	1 1399	7.15	0 254	0.00	0 789	0.00
85-94	1 340	29.41	0 69	0.00	1 159	62.89
Total	3 12647	2.37	0 3282	0.00	3 8171	3.67
MALE						
Age Group						
25-34	0 382	0.00	0 103	0.00	0 142	0.00
35-44	0 980	0.00	0 268	0.00	0 379	0.00
45-54	0 2109	0.00	0 495	0.00	0 779	0.00
55-64	2 2371	0.44	0 483	0.00	0 995	0.00
65-74	0 1683	0.00	0 311	0.00	2 596	33.58
75-84	3 878	34.17	0 61	0.00	0 204	0.00
85-94	0 212	0.00	0 6	0.00	0 11	0.00
Total	6 8616	6.80	0 1728	0.00	2 3107	6.44

TABLE 58

ATISMOO - CURRENT NON SMOKERS
LUNG CANCER (1977-1982)
DISTRIBUTION OF INCIDENT CASES, PERSON YEARS AND INCIDENCE RATES BY AGE GROUP
AND YEARS WORKED WITH A SMOKER

FEMALE Age Group	YEARS WORKED WITH SMOKER					
	NONE		1-10 YEARS		11+ YEARS	
	Cases/ Person years	Rate/ 10,000 PY	Cases/ Person years	Rate/ 10,000 PY	Cases/ Person years	Rate/ 10,000 PY
25-34	0 621	0.00	0 688	0.00	0 165	0.00
35-44	0 1463	0.00	0 1195	0.00	0 583	0.00
45-54	0 2355	0.00	0 1640	0.00	0 1197	0.00
55-64	0 3532	0.00	1 1397	7.18	0 1610	0.00
65-74	1 3379	2.94	0 768	0.00	1 919	10.98
75-84	1 1953	5.12	0 255	0.00	0 234	0.00
85-94	2 439	45.68	0 72	0.00	0 59	0.00
Total	4 13641	2.83	1 5802	1.72	1 4657	2.16
MALE						
Age Group						
25-34	0 313	0.00	0 238	0.00	0 75	0.00
35-44	0 760	0.00	0 695	0.00	0 272	0.00
45-54	0 1797	0.00	0 838	0.00	0 650	0.00
55-64	2 2242	8.92	0 822	0.00	0 785	0.00
65-74	2 1718	11.64	0 440	0.00	0 432	0.00
75-84	1 860	11.63	2 98	204.08	0 186	0.00
85-94	0 193	0.00	0 30	0.00	0 7	0.00
Total	5 7883	8.34	2 3160	6.33	0 2408	0.00

108

2023513054

There were no cases of lung cancer for either sex in the subjects who had lived for 1-10 years with a smoker. For males there were no cases among those who had worked eleven years or more with a smoker.

Four of the thirteen cases occurred among former smokers all of whom smoked more than ten pack years of cigarettes. In a stratified analysis, adjusting for age and sex, the rate ratio of lung cancer among those smoking more than ten pack years compared to those who were non-smokers was 2.81 (95% C.I. 0.78-10.20, $p=0.08$). Consequently, past active smoking exposure was treated as a confounder and controlled for by stratification in additional analyses.

The results of stratified analyses for selected exposure factors controlling for age and subjects past smoking status are presented in Table 5.9 for females and Table 5.10 for males. The differences in the crude RR and adjusted RR indicate some confounding due to past active smoking and/or age differences in the population and therefore the adjusted summary rate ratios are considered as unconfounded by the subjects past smoking status or age. For females who had lived eleven or more years with a smoker as compared to females who had never lived with a smoker, the rate ratio of lung cancer was 1.16 (95% C.I. 0.20-6.61). For females who had worked eleven or more years with a smoker the corresponding rate ratio was 1.47 (95% C.I. 0.15-14.06).

2023513056

110

TABLE 5.B

ATSMOQ - FEMALES
ADJUSTED RATE RATIOS (RR) OF LUNG CANCER (1977-1982)
FOR SELECTED EXPOSURE FACTORS

Factor	Levels	Cases/ Person Years	Rate/ 10 ⁶ PY	Crude RR	Adjusted RR* (95% CI)
Years Lived with Smoker	None 1-10 years 11+ years	3 12818 0 3301 3 8213	0 0 365	Ref. Ref. 1.66	Ref. Ref. 1.18 (0.20-6.61) p = 0.68
Years Worked with Smoker	None 1-10 years 11+ years	4 13861 1 6802 1 4670	289 172 214	Ref. 0.60 0.74	Ref. 1.03 (0.11-10.11) 1.47 (0.15-14.06) p = 0.98
Hours of Oxidant	0-160 161-3000 > 3000	1 6273 4 13424 1 4635	169 298 216	Ref. 1.87 1.38	Ref. 1.73 (0.20-16.25) 0.80 (0.05-14.93) p = 0.80
Hours of TSP	0-100 101-3000 > 3000	1 6328 4 13039 1 6966	188 307 168	Ref. 1.63 0.89	Ref. 1.25 (0.14-11.20) 0.68 (0.04-9.81) p = 0.83
Education	High School College +	3 9213 3 14968	328 200	Ref. 0.61	Ref. 1.22 (0.24-6.22) p = 0.82
Bois Carotene Index	High Medium Low	2 8590 3 7197 1 8228	233 417 121	Ref. 1.78 0.52	Ref. 1.81 (0.30-10.85) 0.35 (0.03-4.29) p = 0.41

*Discrepancies in cases and person years are due to

missing values in exposure factors

* Rate per 1 million person years

Cancer Cases - 6

* Mantel-Haenszel summary rate ratio (RR) adj for age & smoking status

Total Subjects - 4,024

Total Person Years - 24,100

TABLE 5.10

ATISMOX - MALES
ADJUSTED RATE RATIOS (RR) OF LUNG CANCER (1977-1982)
FOR SELECTED EXPOSURE FACTORS

Factor	Levels	* Cases/ Person Years	* Rate/ 10 ⁶ PY	Crude RR	Adjusted RR [†] (95% C.I.)	
Years Lived with Smoker	None	6 8723	673	Ref.	Ref.	
	1-10 years	0 1729	0	0.00	0.00	
	11+ years	2 3123	640	1.12	1.17 (0.21-6.81)	p = 0.64
Years Worked with Smoker	None	5 7996	625	Ref.	Ref.	
	1-10 years	2 3159	633	1.01	1.72 (0.33-9.04)	p = 0.35
	11+ years	0 2420	0	0.00	0.00	
Hours of Oxidant > 10 ppm	0-160	0 3504				
	161-3000	4 7429				
	> 3000	3 2642				
Hours of TSP > 200 ppm	0-100	0 3064				
	101-3000	3 6894				
	> 3000	4 3618				
Education	High School	3 3821	785	Ref.	Ref.	
	College +	3 9692	310	0.39	0.66 (0.17-3.77)	p = 0.62
North Carolina Index	High	1 4114	243	Ref.	Ref.	
	Medium	3 4019	746	3.07	4.01 (0.41-39.46)	
	Low	2 5302	377	1.55	1.98 (0.18-21.73)	p = 0.23

* Discrepancies in cases and person years are due to missing values in exposure factors

* Rate per 1 million person years

* Adjusted RR: Mantel-Haenszel summary rate ratio (RR) adj. for age & smoking status

Total Subjects - 2,261

Cancer Cases - 7

Total Person Years - 13,455

2023513057

Similar results are observed in Table 5.10 for the male cohort. There is a slight increased risk for subjects who lived eleven or more years with a smoker as compared to those who had not lived with a smoker, $RR = 1.17$ (0.21-6.61). For ETS exposure at work the rate ratio of lung cancer in males is 1.72 (95% C.I. 0.33-9.04). Since we have very few cases the conditional maximum likelihood RR estimate and the exact mid probability binomial confidence intervals were also calculated and the results are detailed in Table 5.11. These results are similar to the Mantel-Haenszel stratified analysis.

5.3 SUMMARY

In both populations analyzed there appears to be a positive effect of passive smoking exposure with the outcome of lung cancer. Each of the adjusted measures of effect for all the ETS variables show a positive effect for exposure. However, the magnitude of that effect varies depending on the cohort observed and the particular exposure variable used. For both sexes in the AHSMOG cohort the results indicate that working with a smoker has a greater effect on lung cancer than living with a smoker. However, the results should be interpreted cautiously because of the small number of cases that occurred in both populations. Further discussion of the results are presented in the final chapter.

TABLE E.11

AHSMOG - LUNG CANCER
 * ADJUSTED RATE RATIOS FOR VARYING YEARS
 AND TYPES OF PASSIVE SMOKING EXPOSURE

Passive Smoke Exposure	Females		Males	
	Cases/PY	RR (95% CI)*	Cases/PY	RR (95% CI)*
Years Lived with Smoker				
None	3/12818	Ref.	5/8723	Ref.
1-10	0/3301	0.00	0/1729	0.00
11 +	3/8213	1.22 (0.18-7.89) p = 0.82	2/3123	1.18 (0.15-8.74) p = 0.83
Years Worked with Smoker				
None	4/13861	Ref.	5/7996	Ref.
1-10	1/5802	1.01 (0.04-9.09)	2/3159	1.68 (0.22-8.81)
11 +	1/4670	1.38 (0.05-12.66) p = 0.76	0/2420	0.00 p = 0.55

* Conditional maximum likelihood estimates of RR adjusted for age and past smoking status

- Mid probability (Miettinen) binomial confidence interval